Please write clearly in I	block capitals.		
Centre number		Candidate number	
Surname			
Forename(s)			
Candidate signature			

GCSE PHYSICS

Higher Tier

Paper 1

Wednesday 23 May 2018

Afternoon

Time allowed: 1 hour 45 minutes

Materials

For this paper you must have:

- a ruler
- a scientific calculator
- the Physics Equation Sheet (enclosed).

Instructions

- Use black ink or black ball-point pen.
- Fill in the boxes at the top of this page.
- Answer all questions in the space provided.
- Do all rough work in this book. Cross through any work you do not want to be marked.

• In all calculations, show clearly how you work out your answer.

Information


*

- The maximum mark for this paper is 100.
- The marks for questions are shown in brackets.
- You are expected to use a calculator where appropriate.

• You are reminded of the need for good English and clear presentation in your answers.

For Exami	iner's Use
Question №	lark
1	
2	
3	
4	
5	
6	
7	
8	
9	
10	
TOTAL	





0 1 4	Some carpets have thin copper wires running through them. The student is less likely to	Do not write outside the box
	receive an electric shock after walking on this type of carpet.	
	Suggest why. [2 marks]	
		8

		Do not write
02	A teacher used a Geiger-Muller tube and counter to measure the number of counts in 60 seconds for a radioactive rock.	outside the box
021	The counter recorded 819 counts in 60 seconds. The background radiation count rate was 0.30 counts per second.	
	Calculate the count rate for the rock.	
	[3 marks]	
	Count rate = per second	
02.2	A householder is worried about the radiation emitted by the granite worktop in his kitchen.	
	1 kg of granite has an activity of 1250 Bq. The kitchen worktop has a mass of 180 kg.	
	Calculate the activity of the kitchen worktop in Bq.	
	[2 marks]	
	Activity = Bq	
	Question 2 continues on the next page	
	Turn over ►	

05

02.3	The average total radiation dose per year in the UK is 2.0 millisieverts.			
	Table 1 shows the effects of radiation dose on the human body.			
	Table 1			
	Radiation dose	Effects		
	in millisieverts			
	10 000 Immediate illness; death w	ithin a few weeks		
	1000 Radiation sickness; unlikely t	o cause death		
	100 Lowest dose with evidence of	causing cancer		

The average radiation dose from the granite worktop is 0.003 millisieverts per day.

Explain why the householder should not be concerned about his yearly radiation dose from the granite worktop.

One year is 365 days.


[2 marks]

0 2.4


Bananas are a source of background radiation. Some people think that the unit of radiation dose should be changed from sieverts to Banana Equivalent Dose.

Suggest one reason why the Banana Equivalent Dose may help the public be more aware of radiation risks.

[1 mark]





0 3 2	Describe how the student would obtain the data needed for the investigation.	Do not write outside the box
	Your answer should include a risk assessment for one hazard in the investigation. [6 marks]	
033	Why would switching off the circuit between readings have improved the accuracy of the student's investigation?	
	Tick one box. [1 mark]	
	The charge flow through the wire would not change.	
	The potential difference of the battery would not increase.	
	The power output of the battery would not increase.	
	The temperature of the wire would not change.	




12

Turn over ►








044	Calculate the percentage increase in speed of the cyclist when the power output changes from 200 W to 300 W. [2 marks]	Do not write outside the box
	Percentage increase in speed =	
045	The maximum speed this cyclist can travel on a level road is 14 m/s.	
	How does cycling uphill affect the maximum speed of this cyclist?	
	Explain your answer. [3 marks]	
	Turn over ►	

0 5 1	Complete the se	ntence. Choose answers fro	om the box.		[2 marks]	Do not write outside the box
	charge	potential difference	power	temperature	time	
	The current throu	ugh an ohmic conductor is c		ortional to the aponent, provided		
	that the r <u>emains</u>	constant.				
0 5 2	Figure 7 shows a	a current – potential differe	nce graph foi	a filament lamp.		
	Explain how the across it increase	resistance of a filament lam	ntial difference p changes as		erence [3 marks]	




Do not write outside the A Light Dependent Resistor (LDR) is used to turn on an outside lamp when it gets dark. Part of the circuit is shown in Figure 8. Figure 8 2000 Ω 12V to outside The light intensity decreases. lamp circuit What happens to the potential difference across the LDR and the current in the LDR? [2 marks] Potential difference Current 0 5.4 0 5.5 What is the resistance of the LDR when the potential difference across it is 4 V? Give a reason for your answer. [2 marks] Resistance = Reason

box

0 5 6	Calculate the current through the LDR when the resistance of the LDR is 50)00 Ω.	Do not write outside the box
	Give your answer to 2 significant figures.	[4 marks]	
	Current =	А	14
	Turn over for the next question		
	Ti	urn over 🕨	


0 6	Smoke alarms contain an alpha radiation source and a radiation detector.	Do not wri outside th box
	Figure Part of the inside of a smoke alarm.	
	Figure 9	
	Alpha radiation Smoke	
	Alpha radiation source	
	Plastic casing O O O O Radiation detector Smoke particles	
061	The smoke alarm stays off while alpha radiation reaches the detector. Why does the alarm switch on when smoke particles enter the plastic casing?	
	[1 mark]	
06.2	Why is it safe to use a source of alpha radiation in a house? [1 mark]	

063	The smoke alarm would not work with a radiation source that emits beta or gamma radiation.	Do not write outside the box
	Explain why. [2 marks]	
	Question 6 continues on the next page	
	Turn over ►	
19		



Do not write outside the


box




Explain the ideal properties of a radioactive source for use in medical diagnosis.

[4 marks]

Do not write outside the box





One of the student's results is given below.	Do not write outside the box
pressure = 1.6 × 105 Pa volume = 9.0 cm3 Calculate the volume of the gas when the pressure was 1.8 × 105 Pa.	
The temperature of the gas was constant.	
[3 marks]	

Volume = cm3

0 7.4

25

0 7 5 Figure 14 shows a person using a bicycle pump to inflate a tyre. Figure 14 The internal energy of the air increases as the tyre is inflated. Explain why. Turn over for the next question


Turn over ►

[2 marks]

0 8	Nuclear power stations generate electricity through nuclear fission. Electricity can also be generated by burning shale gas.	Do not write outside the box
081	Shale gas is natural gas trapped in rocks. Shale gas can be extracted by a process called fracking. There is some evidence that fracking causes minor earthquakes. Burning shale gas adds carbon dioxide to the atmosphere.	
	Describe the advantages of nuclear power compared with the use of shale gas to generate electricity. [3 marks]	
082	What is the name of one fuel used in nuclear power stations? [1 mark]	

083	Describe the process of nuclear fission. [4 marks]	Do not write outside the box
		8
	Turn over for the next question	
	Turn over ►	


Do not write outside the box



	The coffee machine heats water from 20 °C to 90 °C.	Do not write outside the box
093		
	The power output of the coffee machine is 2.53 kW.	
	The specific heat capacity of water is 4200 J/kg °C.	
	Calculate the mass of water that the coffee machine can heat in 14 seconds. [5 marks]	
	Mass =kg	
		9
	Turn over for the next question	
	Turn over ►	•

10	Figure 16 shows a wind turbine.	Do not write outside the box
	Figure 16	
101	At a particular wind speed, a volume of 2.3 ×104 m3 of air passes the blades each second.	
	The density of air is 1.2 kg/m3.	
	Calculate the mass of air passing the blades per second.	
	[3 marks]	
	Mass of air per second = kg	
102	The power output of the turbine is directly proportional to the kinetic energy of the air passing the blades each second.	
	Describe the effect on the power output when the wind speed is halved. [3 marks]	

103	At a different wind speed, the wind turbine has a power output of 388 kW. The mass of air passing the wind turbine each second is 13 800 kg. Calculate the speed of the air passing the blades each second. Assume that the process is 100% efficient.	[3 marks]	Do not write outside the box
	Speed of air =	m/s	9
	END OF QUESTIONS		
	-	Turn over ►	



Permission to reproduce all copyright material has been applied for. In some cases, efforts to contact copyright-holders may have been unsuccessful and AQA will be happy to rectify any omissions of acknowledgements. If you have any queries please contact the Copyright Team, AQA, Stag Hill House, Guildford, GU2 7XJ.

Copyright © 2018 AQA and its licensors. All rights reserved.

32