

Mark schemes

Q1.

- | | | |
|-----|--|-----|
| (a) | they form ions with different charges | 1 |
| | they have high melting points | 1 |
| (b) | the (grey) crystals are silver | 1 |
| | the copper ions (produced) are blue
<i>allow the copper nitrate / compound
(produced) is blue</i> | 1 |
| | (because) copper displaces silver | 1 |
| (c) | Level 2: The method would lead to the production of a valid outcome.
The key steps are identified and logically sequenced. | 3-4 |
| | Level 1: The method would not lead to a valid outcome. Some
relevant steps are identified, but links are not made clear. | 1-2 |
| | No relevant content | 0 |
| | Indicative content | |
| | Key steps | |
| | <ul style="list-style-type: none"> • add the metals to (dilute) hydrochloric acid • measure temperature change
or
compare rate of bubbling
or
compare colour of resulting solution | |
| | for copper: | |
| | <ul style="list-style-type: none"> • no reaction • shown by no temperature change
or
shown by no bubbles | |
| | for magnesium and iron: | |
| | <ul style="list-style-type: none"> • magnesium increases in temperature more than iron
or
magnesium bubbles faster than iron
or
magnesium forms a colourless solution and iron forms a | |

coloured solution

Control variables

- same concentration / volume of hydrochloric acid
- same mass / moles of metal
- same particle size of metal
- same temperature (of acid if comparing rate of bubbling)

(d)

$$\frac{(203 \times 30) + (205 \times 70)}{100}$$

or

$$\frac{6090 + 14\,350}{100}$$

= 204.4

ignore units


1

1

[11]

Q2.

(a)



scores 2 marks

allow dots, crosses, circles or e(-) for electrons

1 bonding pair of electrons in each overlap

1

2 non-bonding electrons on nitrogen

do not accept non-bonding electrons on hydrogen

ignore inner shell electrons drawn on nitrogen

1

(b) does not show the shape

or

only two-dimensional


allow is not three-dimensional

- 1
- (c) (ammonia has) small molecules
allow (ammonia has) a simple molecular (structure)
- 1
- (ammonia has) weak intermolecular forces
allow (ammonia has) weak intermolecular bonds
do not accept weak covalent bonds
- 1
- (so) little energy is needed to overcome the intermolecular forces
allow (so) little energy is needed to break the intermolecular bonds
allow (so) little energy is needed to separate the molecules
do not accept references to breaking covalent bonds
- 1
- (d) Cr₂O₃
- 1
- (e)
- an answer of (-)1272 (kJ) scores 3 marks*
- (for bonds broken)
 $((12 \times 391) + (3 \times 498) =) 6186$
- 1
- (for bonds made)
 $((2 \times 945) + (12 \times 464) =) 7458$
- 1
- (overall energy change = $6186 - 7458 =$) (-)1272 (kJ)
allow correct calculation using incorrectly calculated values from step 1 and/or step 2
- 1
- (f)
- allow ecf from part (e)*
- 7458 (kJ) (released in making bonds) is greater than 6186 (kJ) (used in breaking bonds)
 or
 the products have 1272 (kJ) less energy than the reactants
- allow the (overall) energy change is -1272 (kJ)*
- 1
- (so) energy is released (to the surroundings)

*dependent on MP1 being awarded
allow (so) heat is released (to the surroundings)
if no values given, allow 1 mark for more energy released in making bonds than used in breaking bonds*

1

(g)



*scores 2 marks
allow discontinuous lines
ignore arrow heads*

activation energy labelled

1

(overall) energy change labelled

1

[14]

Q3.

(a) chlorine is toxic

*allow carbon monoxide is toxic
allow poisonous for toxic
ignore harmful / deadly / dangerous
allow a poisonous gas is used / produced
allow titanium chloride is corrosive*

1

(b) any one from:

- very exothermic reaction

*allow explosive
allow violent reaction
ignore vigorous reaction
ignore sodium is very reactive*

- produces a corrosive solution
allow caustic for corrosive
ignore alkaline
 - produces hydrogen, which is explosive / flammable
allow flames produced
ignore sodium burns
- (c) argon is unreactive / inert
allow argon will not react (with reactants / products / elements)
- oxygen (from air) would react with sodium / titanium
or
water vapour (from air) would react with sodium / titanium
allow elements / reactants / products for sodium / titanium
- (d) metal chlorides are usually ionic
allow titanium chloride is ionic
- (so)(metal chlorides) are solid at room temperature
or
(so)(metal chlorides) have high melting points
allow titanium chloride for metal chlorides
- (because) they have strong (electrostatic) forces between the ions
ignore strong ionic bonds
- or
(but) must be a small molecule or covalent
allow molecular
- allow alternative approach:*
titanium chloride must be covalent or has small molecules (1)
with weak forces between molecules
do not accept bonds unless intermolecular bonds(1)
(but) metal chlorides are usually ionic (1)
- (e) sodium (atoms) lose electrons
do not accept references to oxygen
- (f) $\text{Na} \rightarrow \text{Na}^+ + \text{e}^-$
do not accept e for e-

1

(g) (Mr of TiCl_4 =) 190

$$\left(\text{moles Na} = \frac{20\,000}{23} =\right) 870 \text{ (mol) }^*$$

1

$$\left(\text{moles TiCl}_4 = \frac{40\,000}{190} =\right) 211 \text{ (mol) }^*$$

1

**allow 1 mark for 0.870 mol Na and
0.211 mol TiCl_4
allow use of incorrectly calculated Mr
from step 1*

either

(sodium is in excess because) 870 mol Na is more than the 844 mol needed

or

(because) 211 mol TiCl_4 is less than the 217.5 mol needed

*the mark is for correct application of the
factor of 4*

*other correct reasoning showing, with
values of moles or mass, an excess of
sodium or insufficient TiCl_4 is
acceptable*

*allow use of incorrect number of moles
from steps 2 and / or 3*

alternative approaches:

approach 1:

(Mr of TiCl_4 =) 190(1)

(40 kg TiCl_4 needs)

$$\frac{40}{190} \times 4 \times 23 \text{ (kg Na) (1)}$$

(=) 19.4 (kg) (1)

so 20 kg is an excess (1)

approach 2:

(Mr of TiCl_4 =) 190(1)

(20 kg Na needs)

$$\frac{20}{4 \times 23} \times 190 \text{ (kg TiCl}_4\text{) (1)}$$

(=) 41.3 (kg) (1)

so 40 kg is not enough (1)

1

- (h) $(\text{actual mass} =) \frac{92.3}{100} \times 13.5$ 1
 or
 $(\text{actual mass} =) 0.923 \times 13.5$ 1

 $= 12.5 \text{ (kg)}$
allow 12 / 12.46 / 12.461 / 12.4605 (kg) 1
an answer 12.5 (kg) scores 2 marks
 [15]

Q4.

- (a) incomplete combustion 1

 (because) insufficient / limited oxygen supply 1
- (b) any two from:
 • carbon monoxide toxic / poisonous
allow description of how carbon monoxide is toxic / poisonous
ignore carbon monoxide is harmful / dangerous / deadly

 • greater public concern / awareness about pollution
ignore comments about the effects of other pollutants
ignore unspecified comments about carbon monoxide pollution

 • more cars so otherwise there would be more carbon monoxide entering atmosphere
 • improved engine technology
 • catalytic converters have been introduced 2
- (c) any one from:
 • (to reduce) health problems
allow (to reduce) specified health problems e.g. breathing difficulties, asthma, lung cancer

 • (to reduce) global dimming
allow (to reduce) the effects of global dimming e.g. reduced light levels
allow (to reduce) smog
allow (to reduce) the formation of particulates


- ignore global warming*
do not accept to reduce soot
- 1
- (d) nitrogen (from atmosphere) reacts with oxygen (from atmosphere)
- 1
- at high temperature (in engine)
ignore heat / hot
- or
with a spark (from spark plug)
- 1
- (e) $2 \text{NO}_2 \rightarrow \text{N}_2 + 2 \text{O}_2$
- allow multiples*
if incorrect, allow N2 for 1 mark
- 2
- (f) any one from:
- acid rain
allow specific effects of acid rain
 - respiratory problems
allow specific respiratory problems e.g. breathing difficulties, asthma
 - carbon monoxide
 - global dimming or smog
- 2
- max 1 mark if global warming mentioned*
- (g) transition metals
- 1
- [12]

Q5.

- (a) in a closed system
- 1
- the rate of the forward and backward reactions are equal
- 1
- (b) concentration increases
- 1
- (because) reaction / equilibrium moves to the left / reactant side
- 1
- (since the) reverse reaction is exothermic
allow (so that) temperature increases
- 1

- (c) becomes blue 1
- (because) reaction / equilibrium moves to the right / product side 1
- (so) concentration of blue cobalt compound increases
allow (so that) concentration of hydrochloric acid decreases 1
- (d) (cobalt has) ions with different charges
allow (cobalt is a) transition metal 1
- (e) Co^{3+} 1
- (f) they allow reactions to reach equilibrium more quickly 1
- they provide a different reaction pathway 1
- (g) $13\text{H}_2 + 6\text{CO} \rightarrow \text{C}_6\text{H}_{14} + 6\text{H}_2\text{O}$
allow multiples 1
- (h) C_8H_{18} 1
- (i) curve below printed curve
do not accept different reactant or product levels 1
- vertical arrow from reactant level to peak of printed curve 1

an answer of:



scores 2 marks

[16]

Q6.

- (a) 13 (protons)

*The answers must be in the correct order.
if no other marks awarded, award 1 mark if number
of protons and electrons are equal*

- 1
- 14 (neutrons) 1
- 13 (electrons) 1
- (b) has three electrons in outer energy level / shell
allow electronic structure is 2.8.3 1
- (c) Level 3 (5–6 marks):
A detailed and coherent comparison is given, which demonstrates a broad knowledge and understanding of the key scientific ideas. The response makes logical links between the points raised and uses sufficient examples to support these links.
- Level 2 (3–4 marks):
A description is given which demonstrates a reasonable knowledge and understanding of the key scientific ideas. Comparisons are made but may not be fully articulated and / or precise.
- Level 1 (1–2 marks):
Simple statements are made which demonstrate a basic knowledge of some of the relevant ideas. The response may fail to make comparisons between the points raised.
- 0 marks:
No relevant content.
Indicative content
Physical
Transition elements
- high melting points
 - high densities
 - strong
 - hard
- Group 1
- low melting points
 - low densities
 - soft
 -
- Chemical
Transition elements
- low reactivity / react slowly (with water or oxygen)
 - used as catalysts
 - ions with different charges
 - coloured compounds
- Group 1
- very reactive / react (quickly) with water / non-metals
 -

- not used as catalysts
- white / colourless compounds
- only forms a +1 ion

6

[10]