Please check the examination details below before entering your candidate information		
Candidate surname	Other names	
Pearson Edexcel Level 1/Level 2 GCSE (9–1)	ntre Number Candidate Number	
Wednesday 10	June 2020	
Morning (Time: 1 hour 10 minutes)	Paper Reference 1SC0/2CH	
Combined Science Paper 5	ze	
	Higher Tier	
You must have: Calculator, ruler	Total Marks	

Instructions

- Use **black** ink or ball-point pen.
- **Fill in the boxes** at the top of this page with your name, centre number and candidate number.
- Answer **all** questions.
- Answer the questions in the spaces provided
 there may be more space than you need.
- Calculators may be used.
- Any diagrams may NOT be accurately drawn, unless otherwise indicated.
- You must **show all your working out** with **your answer clearly identified** at the **end of your solution**.

Information

- The total mark for this paper is 60.
- The marks for each question are shown in brackets
 use this as a quide as to how much time to spend on each question.
- In questions marked with an asterisk (*), marks will be awarded for your ability to structure your answer logically showing how the points that you make are related or follow on from each other where appropriate.
- A periodic table is printed on the back cover of this paper.

Advice

- Read each question carefully before you start to answer it.
- Try to answer every question.
- Check your answers if you have time at the end.

Turn over ▶







Answer ALL questions. Write your answers in the spaces provided.

Some questions must be answered with a cross in a box \boxtimes . If you change your mind about an answer, put a line through the box \boxtimes and then mark your new answer with a cross \boxtimes .

- 1 (a) An atom of potassium has atomic number 19 and mass number 39.
 - (i) Give the electronic configuration of this potassium atom.

(1)

(ii) This potassium atom forms the ion K⁺.

Which row shows the number of protons and the number of neutrons in this potassium ion, K⁺?

(1)

A

В

X C

X D

number of protons	number of neutrons
19	19
19	20
20	19
20	20

(b) Potassium and caesium are in the same group of the periodic table.

Explain, in terms of electrons, why potassium and caesium are in the same group.

(2)

(c) Fluorine boils at -188 °C.

There are forces between fluorine molecules.

Explain, in terms of these forces, why the boiling point of fluorine is low.

(2)





(d) Potassium reacts with fluorine to form potassium fluoride. Potassium fluoride is a solid.

Complete the balanced equation for this reaction and add the state symbols.

(3)

.....K (......) +
$$F_2(g)$$
 \rightarrow KF (......)

(Total for Question 1 = 9 marks)

2 Calcium carbonate reacts with dilute hydrochloric acid to produce carbon dioxide gas.

The rate of reaction between calcium carbonate and dilute hydrochloric acid at room temperature was investigated.

(a) The investigation was carried out with different sized calcium carbonate pieces.

The mass of calcium carbonate and all other conditions were kept the same.

The results are shown in Figure 1.

size of calcium carbonate pieces used	volume of carbon dioxide gas produced in five minutes in cm ³
large	16
small	48
powder	90

Figure 1

State, using the information in Figure 1, the effect of the surface area of the calcium carbonate on the rate of this reaction.	(1)
(b) The calcium carbonate powder produced $90\mathrm{cm^3}$ of carbon dioxide in five minutes Calculate the average rate of reaction in $\mathrm{cm^3s^{-1}}$.	(3)
average rate of reaction –	3 s:

(c) The experiments were repeated at a higher temperature. The rate of reaction for each experiment increased.	
Explain, in terms of particles, why the rate of reaction increased when the temperature was increased.	
	(3)
(Total for Question 2 =	7 marks)

3 (a) Sodium thiosulfate solution, Na₂S₂O₃, reacts with dilute hydrochloric acid.

$$Na_2S_2O_3(aq) + 2HCl(aq) \rightarrow 2NaCl(aq) + H_2O(l) + SO_2(g) + S(s)$$

(i) When dilute hydrochloric acid is mixed with sodium thiosulfate solution, the mixture turns cloudy.

Explain why the mixture turns cloudy.

(2)

(ii) In an investigation, different concentrations of hydrochloric acid are reacted with sodium thiosulfate solution.

The mixture goes cloudy at different rates.

Describe how the rate at which the mixture goes cloudy can be measured.

(3)

(iii) You are provided with some dilute hydrochloric acid which has a concentration of $50\,\mathrm{g\,dm^{-3}}$.

For this experiment, dilute hydrochloric acid with a concentration of $20\,\mathrm{g}\,\mathrm{dm}^{-3}$ is required.

How much water must be added to 100 cm³ of 50 g dm⁻³ hydrochloric acid to make dilute hydrochloric acid with a concentration of 20 g dm⁻³?

(1)

- B 150 cm³



(b) Sodium iodide solution is colourless.	
When a solution of bromine is added to sodium iodide solution, a reaction occurs.	
$2NaI + Br_2 \rightarrow 2NaBr + I_2$	
(i) The mixture turns brown.	
Give the name of the substance causing the brown colour.	
	(1)
(ii) Explain which substance has been reduced in this reaction.	
(II) Explain which substance has been reduced in this reaction.	(2)
	,
(Total for Question 3 – 9 ma	rks)

4 (a) Air contains several gaseous elements.

Which of these shows the three most common gaseous elements in air, listed in order from the most common to the least common?



- A oxygen, chlorine, nitrogen
- **B** nitrogen, oxygen, hydrogen
- C oxygen, nitrogen, helium
- D nitrogen, oxygen, argon
- (b) The density of a gas can be found using the equation

$$density = \frac{mass}{volume}$$

A student carried out an experiment to find the density of argon.

The mass of a stopper and flask, containing no gas, was known. The flask was completely filled with argon and its mass measured.

Figure 2 shows the results the student wrote down.

mass of stopper and flask in g	78.639
mass of stopper and flask full of argon in g	79.120
volume of flask in cm ³	250.0

Figure 2

(i)	Use the results	to calculate	the density o	of argon in $g cm^{-3}$.
-----	-----------------	--------------	---------------	---------------------------

- 4	10	9	1
- (J	-1
٠,	- 4	ь	

density of argon -	q	cm-3

(ii) The flask used for the experiment is shown in Figure 3. The flask holds 250.0 cm³ when filled up to the line.




Figure 3

There is an error in the volume the student has used in the calculation. This would give an incorrect value for the density of argon.

Identify this error and state what should be done to correct it.

what should be done to correct it

(c) Four of the noble gases are argon, helium, krypton and neon.

Give these gases in order of increasing density.

(2)

(2)

(2)

(d) Much of the carbon dioxide present in the Earth's early atmosphere dissolved into the oceans.

This led to the formation of compounds including calcium carbonate, CaCO₃.

Some of the calcium carbonate reacted with magnesium ions to form dolomite, CaMg(CO₃)₂.

Complete the **ionic** equation for the reaction of calcium carbonate with magnesium ions.

.....CaCO $_3$ + \rightarrow CaMg(CO $_3$) $_2$ + Ca $^{2+}$ dolomite

(e) **P** and **Q** are both mixtures of gases.

One has the same composition as the early atmosphere and the other has the same composition as the current atmosphere.

Tests are carried out on gas mixtures P and Q.

The test for carbon dioxide is to bubble the gas into limewater; if carbon dioxide is present calcium carbonate is formed.

The results of the tests are shown in Figure 4.

test	result with gas mixture P	result with gas mixture Q
bubble gas into limewater	white precipitate forms after 4 minutes	white precipitate forms after 10 seconds
place burning splint into gas mixture	splint continues to burn	splint immediately goes out

Figure 4

(Total for Question 4 = 11 ma	rks)
	(2)
explain, using the data in Figure 4, which gas mixture represents the early atmosp	nere.



BLANK PAGE



- **5** The first four elements in group 1 are lithium, sodium, potassium and rubidium.
 - (a) Rubidium reacts with water to form rubidium hydroxide and hydrogen.

$$2Rb(s) + 2H_2O(l) \rightarrow 2RbOH(aq) + H_2(g)$$

(i) Predict what you would **see** when a small piece of rubidium is placed in a large volume of water.

(3)

(ii) Why is rubidium more reactive than potassium?

(1)

- A the metallic bonds in rubidium are weaker than those in potassium
- B rubidium is a softer metal than potassium
- the outer electron of a rubidium atom is further from the nucleus than potassium's
- **D** rubidium has a more exothermic reaction with water than potassium does
- (iii) 8.5 g of rubidium are reacted completely with water.

The reaction makes a solution of rubidium hydroxide.

The volume of this solution is 2.5 dm³.

Calculate the concentration of the rubidium hydroxide solution in g dm⁻³.

(relative atomic mass: Rb = 85; relative formula mass: RbOH = 102)

(4)

 $concentration = \dots g dm^{-3}$

(b) An example of an endothermic reaction is the reaction between rubidium hydroxic and ammonium carbonate, $(NH_4)_2CO_3$.	de
This reaction forms rubidium carbonate, Rb ₂ CO ₃ , ammonia and one other product.	
Write the balanced equation for this reaction.	(3)
(Total for Question 5 = 11 ma	r k s)

6 (a) An impure hydrocarbon fuel is burned in the apparatus in Figure 5.




Figure 5

When the fuel is burned

- the funnel becomes hot
- a colourless liquid forms in tube A
- the indicator in tube **B** changes colour to show an acidic gas.

Explain these observations.

(3)

(b) The energies of some bonds are shown in Figure 6.

bond	bond energy in kJ mol ⁻¹
С—Н	435
0=0	496
c=o	805
Н—О	463

Figure 6

Methane burns in oxygen to form carbon dioxide and water.

The equation shows the structures of the molecules.

Calculate the energy change, in kJ mol⁻¹, for this reaction.

energy change =kJ mol⁻¹

(4)

*(c) Petrol and diesel are used as fuels for cars.

The emissions from three similar sized cars were investigated.

The first car was the oldest, had no catalytic converter and used petrol.

The other two cars were only a few years old.

One of these was fitted with a catalytic converter and used petrol and the other car used diesel.

Figure 7 shows the emissions in grams for each kilometre travelled by these three cars.

	emissions in g km ⁻¹			
	carbon monoxide	nitrogen oxides	carbon dioxide	carbon particulates
car with no catalytic converter using petrol	1.60	0.09	180	0.00
car with catalytic converter using petrol	0.67	0.02	180	0.00
car using diesel	0.05	0.19	130	0.02

Figure 7

Discuss and compare the impact on the environment of the emissions from these three cars using the information from Figure 7.			2	
	-	-		(6)

TOTAL FOR PAPER = 60 MARKS
(Total for Question 6 = 13 marks)



BLANK PAGE

BLANK PAGE



The periodic table of the elements

0 4 He helium 2	20 Ne neon 10	40 Ar argon 18	84 Kr krypton 36	131 Xe xenon 54	[222] Rn radon 86
7	19 F fluorine 9	35.5 CI chlorine 17	80 Br bromine 35	127 	[210] At astatine 85
9	16 O oxygen 8	32 S sulfur 16	79 Selenium 34	128 Te tellurium 52	[209] Po polonium 84
2	14 N nitrogen 7	31 P phosphorus 15	75 As arsenic 33	122 Sb antimony 51	209 Bi bismuth 83
4	12 carbon 6	28 Si silicon 14	73 Ge germanium 32	119 Sn tin 50	207 Pb lead 82
ဇ	11 boron 5	27 Al aluminium 13	70 Ga gallium 31	115 In indium 49	204 TI thallium 81
'			65 Zn zinc 30	112 Cd cadmium 48	201 Hg mercury 80
			63.5 Cu copper 29	108 Ag silver 47	197 Au gold 79
			59 nickel 28	106 Pd palladium 46	195 Pt platinum 78
			59 Co cobalt 27	103 Rh rhodium 45	192 Ir iridium 77
1 hydrogen			56 iron 26	Ru ruthenium 44	190 Os osmium 76
			55 Mn manganese 25	[98] Tc technetium 43	186 Re rhenium 75
relative atomic mass atomic symbol number atomic (proton) number		52 Cr	96 Mo molybdenum 42	184 W tungsten 74	
		51 V vanadium 23	93 Nb niobium 41	181 Ta tantalum 73	
	relativ ato atomic		48 Ti titanium 22	91 Zr zirconium 40	178 Hf hafnium 72
·			45 Sc scandium 21	89 Y yttrium 39	139 La* lanthanum 57
2	9 Be beryllium 4	24 Mg magnesium 12	40 Ca calcium 20	88 Sr strontium 38	137 Ba barium 56
-	7 Li lifhium 3	23 Na sodium 11	39 K potassium 19	85 Rb rubidium 37	133 Cs caesium 55

^{*} The elements with atomic numbers from 58 to 71 are omitted from this part of the periodic table.

The relative atomic masses of copper and chlorine have not been rounded to the nearest whole number.

