Please check the examination detail	ils bel	ow before ente	ering your candidate information
Candidate surname			Other names
Pearson Edexcel Level 1/Level 2 GCSE (9–1)	Cen	itre Number	Candidate Number
Time 1 hour 10 minutes		Paper reference	1SC0/2CF
Combined Scient PAPER 5 Foundation Tier	nc	e	
You must have: Calculator, ruler			Total Marks

Instructions

- Use **black** ink or ball-point pen.
- **Fill in the boxes** at the top of this page with your name, centre number and candidate number.
- Answer all questions.
- Answer the questions in the spaces provided
 - there may be more space than you need.
- Calculators may be used.
- Any diagrams may NOT be accurately drawn, unless otherwise indicated.
- You must show all your working out with your answer clearly identified at the end of your solution.

Information

- The total mark for this paper is 60.
- The marks for each question are shown in brackets
 - use this as a guide as to how much time to spend on each question.
- In questions marked with an **asterisk** (*), marks will be awarded for your ability to structure your answer logically showing how the points that you make are related or follow on from each other where appropriate.
- There is a periodic table on the back cover of the paper.

Advice

- Read each question carefully before you start to answer it.
- Try to answer every question.
- Check your answers if you have time at the end.
- Good luck with your examination.

Turn over ▶







(1)

Answer ALL questions. Write your answers in the spaces provided.

Some questions must be answered with a cross in a box \boxtimes . If you change your mind about an answer, put a line through the box \boxtimes and then mark your new answer with a cross \boxtimes .

				mark your new answer with a cross $oxtimes$.	
1	Mag	gnesiu	ım re	eacts with dilute sulfuric acid to form magnesium sulfate and hydrogen	gas.
	A st	udent	t wa	nts to find out the effect of temperature on the rate of this reaction.	
	The student used the following method.				
	:	step 1	1 pc	our 25 cm³ of dilute sulfuric acid into a conical flask	
		-		arm the acid until its temperature is 30°C	
		•		d a piece of magnesium to the acid art a stopwatch	
		•		nit until the reaction has finished	
		-		op the stopwatch	
		-	,	peat steps 1–6 but at 50 °C.	
	(a) The student kept the volume of sulfuric acid the same when they repeated the method at 50°C.				
	9	State	two	other variables that should be kept the same.	
				·	(2)
1					
2					
			•		. 12
	(b) \	Which	n pie	ce of equipment can be used to find the volume of 25 cm ³ of sulfuric ac	(1)
		X	A	balance	
		X	В	measuring cylinder	
		X	C	ruler	
		×	D	thermometer	
	(c) :	State	how	the student will know that the reaction has finished.	
					(1)
	(d) ⁻	The re	eacti	on at 50°C was faster than the reaction at 30°C.	
				reason, in terms of particles, why the reaction at 50°C was faster than on at 30°C.	



(Total for Question 1 = 7	marks)
average rate of reaction =	cm³ s ⁻¹
reaction.	(2)
Calculate the average rate of reaction, in cm ³ s ⁻¹ , for the first 60 seconds of the	
(e) At 50°C, 15.0 cm³ of gas was produced during the first 60 seconds of the reacti	on.

- **2** This question is about the noble gases.
 - (a) (i) State, in terms of outer shell electrons, why the noble gases are unreactive.

(1)

(ii) Figure 1 shows an airship, filled with helium, floating above the ground.




Figure 1

Helium, hydrogen and krypton are gases. Figure 2 shows the reactivity and density, at room temperature and pressure, of helium, hydrogen and krypton.

gas	reactivity	density in g cm ⁻³
helium	unreactive	0.00018
hydrogen	very reactive	0.00009
krypton	unreactive	0.00380

Figure 2

The density of air is $0.001225 \,\mathrm{g}\,\mathrm{cm}^{-3}$.

Helium is used in airships.

Explain why hydrogen and why krypton are **not** used in airships.

(b) Mendeleev produced one of the earliest periodic tables.

State why he could **not** include any of the noble gases in his periodic table.

(1)


(c) Figure 3 shows the boiling points of some of the noble gases.

noble gas	boiling point in °C
helium	-269
neon	-246
argon	-186
krypton	-153
xenon	

Figure 3

(i) Complete the bar chart to show the boiling points of neon and argon.

(2)



(ii) Predict the boiling point of xenon.

(1)

boiling point of xenon =°C

(Total for Question 2 = 8 marks)




•			ne, bromine and iodine are elements in group 7 of the periodic table. e name given to the group 7 elements.	(1)
••••			ne other element that is in group 7. periodic table on the back of this exam paper to help you.	(1)
••••	(iii) Whic	ch el	lement is liquid at room temperature and pressure?	(4)
	\times	A	fluorine	(1)
	\boxtimes	В	chlorine	
	\boxtimes	C	bromine	
	\boxtimes	D	iodine	
	(iv) Whic	ch el	lement is dark grey in colour at room temperature and pressure?	(1)
	\boxtimes	A	fluorine	(1)
	\boxtimes	В	chlorine	
	\boxtimes	C	bromine	
	\times	D	iodine	
			ith chlorine to form tin chloride. tin chloride contains 1.19 g of tin and 1.42 g of chlorine.	
	Calculat	e th	e empirical formula of this tin chloride.	
	(relative	ato	mic masses: Cl = 35.5, Sn = 119.0)	
	You mus	st sh	ow your working.	(3)
				(-)
			empirical formula of this tin chloride =	



(c) Tin also reacts with fluorine.

The reaction between fluorine and tin is much more vigorous than the reaction between chlorine and tin.

Figure 4 shows the electronic configurations of fluorine and chlorine.






Figure 4

Explain, in terms of their electronic configurations, why fluorine reacts with tin more vigorously than chlorine reacts with tin.

(2)	

(Total for Question 3 = 9 marks)



BLANK PAGE

4 Figure 5 shows a sample of hydrogen peroxide solution decomposing to form water and oxygen gas.




Figure 5

(a) (i) Write the word equation for hydrogen peroxide solution decomposing.

(1)


(ii) In this reaction hydrogen peroxide is a solution, water is a liquid and oxygen is a gas.


Draw one straight line from each substance to its correct state symbol.

(2)

substance

hydrogen peroxide solution






state symbol







(b) Describe the test to show the gas produced is oxygen.

(2)



(2)

(c) Figure 6 shows the electron arrangement for an atom of hydrogen and an atom of oxygen.




Figure 6

Complete the dot and cross diagram in Figure 7 for a molecule of water, $\rm H_2O$. Draw outer shell electrons only.




Figure 7

(d) Liver contains the enzyme catalase.

A piece of liver was added to another sample of hydrogen peroxide solution. Figure 8 shows the results.





Figure 8





5 (a) The concentration of a solution can be calculated using the equation

concentration of solution =
$$\frac{\text{mass of solid}}{\text{volume of solution}}$$


A student dissolved 9.25 g of ammonium chloride in water and made up the solution to a volume of 200 cm³.

Use the equation to calculate the concentration of this solution in g dm⁻³.

(2)

concentration =g dm⁻³

(b) Dissolving ammonium chloride in water is an endothermic process. Figure 10 shows part of the reaction profile for this process.



progress of reaction

Figure 10


(i) Explain how Figure 10 shows that dissolving ammonium chloride in water is an endothermic process.

(2)

(ii) Complete the reaction profile in Figure 10 and label the activation energy.

(2)

(c) A student used the equipment in Figure 11 to investigate whether electricity can pass through solid ammonium chloride and through ammonium chloride solution.






Figure 11

If an electrical current flows in the circuit, the lamp will light up.

Figure 12 shows the results of the investigation.

substance	lamp
solid ammonium chloride	did not light up
ammonium chloride solution	lit up brightly

Figure 12

Explain the resu	Its of the	investigation.
------------------	------------	----------------

(3)

- (d) Ammonia gas is toxic.
 - (i) Which symbol should be placed on a container of a toxic gas?

(1)









(ii) Give **one** safety precaution that should be taken when working with toxic gases in the laboratory.

(1)

(Total for Question 5 = 11 marks)

(a	Diesel oil is a mixture of hydrocarbons that can be obtained from crude oil. a) State the name of the process used to separate diesel oil from crude oil.						
				(1)			
(Diesel oil contains alkanes. These alkanes are part of an homologous series.					
	Wł	Which statement about compounds in this homologous series is true?					
	X	A	they have the same chemical formula	(1)			
	X	В	they have the same empirical formula				
	X	C	they have the same general formula				
	\times	D	they have the same molecular formula				
		When fuels such as diesel oil are burned, the high temperatures produced can cause nitrogen and oxygen in the air to form the pollutant nitrogen dioxide.					
	Co	mple	ete the balanced equation for the reaction.	(2)			
				(2)			
	$N_2 + 2O_2 \rightarrow \dots$						
	(d) Explain how the greenhouse effect is caused by the gases produced by the complete combustion of diesel oil.						
				(3)			
•••							

*(e)	Diesel oil can contain impurities of sulfur. Burning diesel oil containing impurities of sulfur can result in acid rain. Acid rain is harmful to the environment.							
	Explain how acid rain is formed and the harm it can do.	(6)						





TOTAL FOR PAPER = 60 MARKS

BLANK PAGE



BLANK PAGE



The periodic table of the elements

0	4 He helium 2	20 Ne neon 10	40 Ar argon 18	84 Kr krypton 36	131 Xe xenon 54	[222] Rn radon 86
7		19 F fluorine 9	35.5 CI chlorine 17	80 Br bromine 35	127 	[210] At astatine 85
9		16 O oxygen 8	32 S sulfur 16	79 Se selenium 34	128 Te tellurium 52	[209] Po polonium 84
2		14 N nitrogen 7	31 P phosphorus 15	75 As arsenic 33	122 Sb antimony 51	209 Bi bismuth 83
4		12 C carbon 6	28 Silicon 14	73 Ge germanium	119 Sn th 50	207 Pb lead 82
3		11 B boron 5	27 AI aluminium 13	70 Ga gallium 31	115 In indium 49	204 T thallium 81
	·			65 Zn zinc 30	112 Cd cadmium 48	201 Hg mercury 80
				63.5 Cu copper 29	108 Ag silver 47	197 Au gold 79
				59 Ni nickel 28	106 Pd palladium 46	195 Pt platinum 78
				59 Co cobalt 27	103 Rh rhodium 45	192 Ir iridium 77
	1 hydrogen 1				Ru ruthenium 44	190 Os osmium 76
•				55 Mn manganese 25	[98] Tc technetium 43	186 Re rhenium 75
		mass bol lumber		52 Cr chromium 24	96 Mo molybdenum 42	184 W tungsten 74
	Key	relative atomic mass atomic symbol _{name} atomic (proton) number		51 V vanadium 23	93 Nb niobium 41	181 Ta tantalum 73
				48 Ti titanium 22	91 Zr zirconium 40	178 Hf hafnium 72
			-	45 Sc scandium 21	89 Y yttrium 39	139 La* lanthanum 57
2		9 Be beryllium 4	24 Mg magnesium 12	40 Ca calcium 20	88 Sr strontium 38	137 Ba banum 56
_		7 Li lithium 3	23 Na sodium 11	39 K potassium	85 Rb rubidium 37	133 Cs caesium 55

^{*} The elements with atomic numbers from 58 to 71 are omitted from this part of the periodic table.

The relative atomic masses of copper and chlorine have not been rounded to the nearest whole number.